

# PRIMAP-hist v1.2: updated figures

Johannes Gütschow<sup>\*1</sup>, M. Louise Jeffery<sup>1</sup>, Robert Gieseke<sup>1</sup>, and  
Ronja Gebel<sup>1</sup>

<sup>1</sup>Potsdam Institute for Climate Impact Research, Telegraphenberg  
A 31, 14473 Potsdam, Germany

February 2, 2018

This document contains updates of the figures from Gütschow et al. [2016]  
using PRIMAP-hist v1.2 data.

## References

Johannes Gütschow, M. Louise Jeffery, Robert Gieseke, Ronja Gebel, David Stevens, Mario Krapp, and Marcia Rocha. The PRIMAP-hist national historical emissions time series. *Earth System Science Data*, 8(2):571–603, nov 2016. ISSN 1866-3516. doi: 10.5194/essd-8-571-2016. URL <http://www.earth-syst-sci-data.net/8/571/2016/>.

---

<sup>\*</sup>johannes.guetschow@pik-potsdam.de




Figure 1: Coverage of years and countries in the sources used for the PRIMAP-hist dataset. The color indicates the country group covered or the regional resolution, while the intensities indicates the fraction of countries in the group covered by the source in each year. The fraction is taken over all gases and categories which can be seen in the CDIAC time series where the flaring time series only starts in 1950. RCP time series for CH<sub>4</sub> end in 2000 leading to the lower coverage after year 2000.




Figure 2: Example for the work of the composite source generator: the creation of the category 1A, CO<sub>2</sub> pathway for South Korea. The buildup starts with the UNFCCC source as there is no CRF data for South Korea. Extrapolation is not needed in this case, so the step is omitted from the figure.




Figure 3: Emissions by sector for major emitters and the world. Where land use emissions are negative, the stacked emissions of the other sectors start at this negative value. International shipping and aviation emissions are not included.




Figure 4: Emissions by gas for major emitters and the world. International shipping and aviation emissions are not included.




Figure 5: Comparison of the PRIMAP-hist dataset with both individual sources and composite datasets for aggregate Kyoto gases and the main IPCC 1996 categories. Grey shaded areas show the indicative uncertainty range from Table 10 of Gütschow et al. [2016] applied to the PRIMAP-hist dataset. Where different uncertainty estimates exist the average value is used. International shipping and aviation emissions are not included.




Figure 6: Comparison of the PRIMAP-hist dataset with both individual sources and composite datasets for different gases. Grey shaded areas show the indicative uncertainty range from Table 10 of Gütschow et al. [2016] applied to the PRIMAP-hist dataset. Where different uncertainty estimates exist the average value is used. International shipping and aviation emissions are not included.